

University of Dundee

BACE1 activity impairs neuronal glucose oxidation

Findlay, John A.; Hamilton, David L.; Ashford, Michael L.

Published in:
Frontiers in Cellular Neuroscience

DOI:
[10.3389/fncel.2015.00382](https://doi.org/10.3389/fncel.2015.00382)

Publication date:
2015

Licence:
CC BY

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Findlay, J. A., Hamilton, D. L., & Ashford, M. L. (2015). BACE1 activity impairs neuronal glucose oxidation: rescue by beta-hydroxybutyrate and lipoic acid. *Frontiers in Cellular Neuroscience*, 9, [382].
<https://doi.org/10.3389/fncel.2015.00382>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Bace1 activity impairs neuronal glucose metabolism: rescue by beta-hydroxybutyrate and lipoic acid

John A. Findlay¹, David L. Hamilton¹, Michael L. Ashford^{1*}

¹Division of Cardiovascular & Diabetes Medicine, University of Dundee, United Kingdom

Submitted to Journal:
Frontiers in Cellular Neuroscience

Article type:
Original Research Article

Manuscript ID:
160778

Received on:
14 Jul 2015

Revised on:
28 Aug 2015

Frontiers website link:
www.frontiersin.org

BACE1 activity impairs neuronal glucose oxidation: rescue by beta-hydroxybutyrate and lipoic acid

John A Findlay, D Lee Hamilton, Michael LJ Ashford

Division of Cardiovascular and Diabetes Medicine, School of Medicine, Ninewells Hospital & Medical School, University of Dundee, Dundee, Scotland, UK

Correspondence: Michael LJ Ashford, Division of Cardiovascular and Diabetes Medicine, School of Medicine, Ninewells Hospital & Medical School, University of Dundee, Dundee, DD1 9SY, Scotland, UK. Phone +44 1382 383095, m.l.j.ashford@dundee.ac.uk

Keywords: glucose metabolism, BACE1, amyloid, pyruvate dehydrogenase, mitochondria, alpha lipoic acid

Running Title: BACE1 inhibits neuronal glucose oxidation

Word count: 4681

Figure Count: 6

Abstract

Glucose hypometabolism and impaired mitochondrial function in neurons have been suggested to play early and perhaps causative roles in Alzheimer's disease (AD) pathogenesis. Activity of the aspartic acid protease, beta-site amyloid precursor protein (APP) cleaving enzyme 1 (BACE1), responsible for beta amyloid peptide generation, has recently been demonstrated to modify glucose metabolism. We therefore examined, using a human neuroblastoma (SH-SY5Y) cell line, whether increased BACE1 activity is responsible for a reduction in cellular glucose metabolism. Overexpression of active BACE1, but not a protease-dead mutant BACE1, protein in SH-SY5Y cells reduced glucose oxidation and the basal oxygen consumption rate, which was associated with a compensatory increase in glycolysis. Increased BACE1 activity had no effect on the mitochondrial electron transfer process but was found to diminish substrate delivery to the mitochondria by inhibition of key mitochondrial decarboxylation reaction enzymes. This BACE1 activity-dependent deficit in glucose oxidation was alleviated by the presence of beta hydroxybutyrate or α -lipoic acid. Consequently our data indicate that raised cellular BACE1 activity drives reduced glucose oxidation in a human neuronal cell line through impairments in the activity of specific tricarboxylic acid cycle enzymes. Because this bioenergetic deficit is recoverable by nutraceutical compounds we suggest that such agents, perhaps in conjunction with BACE1

inhibitors, may be an effective therapeutic strategy in the early-stage management or treatment of AD.

Introduction

Alzheimer's disease (AD) is an age-related neurodegenerative disease, with the vast majority of cases described as sporadic and not currently linked with a specific gene defect. There are approximately 44 million people living with AD in the world, and this number is predicted to increase dramatically over the next three decades, severely impacting on healthcare systems and the socio-economic environment. Presently, we have a limited understanding of the early events in sporadic AD disease aetiology. While the progression of AD is closely associated with dysfunctional tau protein and cholinergic deficits, it is generally observed by biomarker analysis that changes in the hallmark amyloid plaque pathology precedes these other pathogenic drivers [1-4]. There has also been increasing interest in the role of brain mitochondrial dysfunction and reduced metabolic activity [5-7].

It has long been recognised that glucose is the predominant substrate utilised by the adult brain under physiological conditions [8]. Indeed, while only constituting 2% to body weight, the brain consumes around 20 and 25% of the total body oxygen consumption and glucose respectively [9]. Furthermore, the respiratory quotient of the brain is almost exactly 1, indicating near universal carbohydrate metabolism [10]. Although glucose is the dominant substrate for brain metabolic activity, alternative substrates, such as glycogen and amino acids can play a role in central metabolism. However due to limited supply and storage capacity, this is thought to be relatively minor under physiological conditions [11]. Glucose metabolism in the brain is tightly coupled to the generation of intracellular adenosine triphosphate (ATP), largely to support neurotransmitter production and release. Consequently, this reliance on glucose coupled with a high-energy consumption makes the brain, and synaptic transmission in particular, vulnerable to events, which lead to diminished metabolism.

Brain hypometabolism is a universal change observed during Alzheimer's disease (AD) progression and recently, it has been proposed to have a major causative role in AD pathogenesis [12,13]. Through the use of neuroimaging techniques, impaired brain glucose metabolism has been demonstrated to occur: before atrophy in autosomal AD cases, during the progression towards non-familial AD and in individuals at high risk of AD (i.e. APOE ϵ 4 carriers) prior to symptom manifestation [14-18]. This reduced brain glucose metabolism has been postulated to result from impaired mitochondrial functioning [19-21], with similar changes observed in a variety of transgenic mouse models of AD [7,22-24]. Recent evidence suggests a putative link between neuronal glucose metabolism and the presence of beta amyloid ($A\beta$) peptides, considered by many to be the primary pathogenic driver of AD. $A\beta$ peptides derive from increased cleavage of amyloid precursor protein (APP), which is ubiquitously expressed in tissues, with highest levels in the brain. APP is cleaved by two competing enzyme processes; an alpha secretase pathway, resulting in the soluble cleaved product sAPP α and no $A\beta$ production and a beta secretase pathway, which releases sAPP β and following additional cleavage of the remaining protein by γ -secretase, releases $A\beta$. $A\beta$ accumulates in the mitochondria of AD patients and AD mouse models, prior to the appearance of amyloid deposits [25-27]. Indeed, exogenous $A\beta$ has been reported to decrease

mitochondrial respiratory chain function and the activity of various mitochondrial dehydrogenase enzymes [28-31], indicative of an A β -mediated bioenergetic deficit in cells.

Additionally, it has been shown that the hypometabolic state associated with AD may be driven, at least in part, by a region-specific shift in neuronal metabolism towards aerobic glycolysis (reduced oxidative metabolism in the presence of adequate oxygen supply; [32]). This switch in glucose metabolism was shown to closely correlate with A β deposition and later vulnerability to cell death during progression towards AD [33]. These findings clearly implicate the modulation of APP processing as a driver of the altered metabolic state observed in early, pre-AD states. A key enzyme driving excess A β production, as observed in AD, is the aspartyl protease, β -site APP cleaving enzyme 1 (BACE1). BACE1 was initially characterised as the enzyme controlling the rate-limiting step in A β generation [34-36]. More recently however, it has also been proposed to play a role in glucose metabolism with whole body knock out of BACE1 resulting in improved insulin sensitivity and glucose homeostasis [37]. Furthermore, genetic and pharmacological manipulation of APP processing can directly alter glucose uptake and metabolism in the C2C12 skeletal muscle cell line [38]. Importantly, BACE1 is a stress-sensitive protease, with oxidative, hypoxic, inflammatory and metabolic stress (all associated with AD initiation and/or progression) demonstrated to increase BACE1 levels and activity, causing APP processing to shift from the physiologically predominant alpha-secretase, to the beta-secretase, pathway and increasing A β levels. Indeed, recent work has demonstrated a role for oxidative, lipid and metabolic stressors in regulating BACE1 gene and protein expression as well as activity [39-43]. Furthermore, alterations in its expression result from changes in micro RNA (miR) regulation of the 5' untranslated region (UTR) of BACE1 have been demonstrated in sporadic AD cases [44-46]. Consequently, chronic stress events and altered translational regulation may in turn culminate in the consistent reports that BACE1 mRNA, protein and activity are elevated in AD brains [47-53].

Given that A β directly impairs mitochondrial enzyme function and that AD is associated with impaired glucose metabolism we hypothesised that manipulating APP processing through BACE1 overexpression in SH-SY5Y neuroblastoma cells would phenocopy the defects in glucose metabolism at the cellular level, allowing us to explore in more detail this initial and potentially causative change in AD progression.

Materials and Methods

Cell culture

SH-SY5Y cells were cultured under aseptic conditions and maintained in a humidified atmosphere of 95% air and 5% CO₂. Cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) F-12 media (Gibco Life Technologies, Paisley, UK) supplemented with 10% fetal bovine serum (Sera Laboratories, West Sussex, UK), 4 mM L-glutamine (Gibco) and 2% Penicillin streptomycin (100 units/ml; Gibco). SH-SY5Y cells were transfected with 12 μ g DNA of pcDNA3.1 containing empty vector (SH-SY5Y_{EV}), full-length human BACE1 (SH-SY5Y_{B1}), or BACE1 active site mutant (SH-SY5Y_{mB1}) using Lipofectamine 2000 (Invitrogen Life Technologies, Paisley, UK). Stable cells were selected and lines maintained using 1 and 0.5 mg/ml G418 sulphate respectively (Sigma-Aldrich, Gillingham, UK) as selection antibiotic. A minimum of 2 independently generated stable cell lines, with

concurrently produced EV controls, were used for these studies. Prior to some glucose oxidation assays, cells were treated overnight, as indicated, with the pyruvate dehydrogenase inhibitor, dichloroacetate (DCA; Sigma-Aldrich) or growth media supplemented with α -lipoic acid (Sigma-Aldrich) for 48 hours.

Cloning

Full length myc-his tagged human BACE1 in pcDNA3.1 was obtained from GlaxoSmithKline (GSK; Harlow, UK) and mBACE1 (a kind gift from Professor Wolfe (Brigham and Women's Hospital, Boston)) was sub-cloned into pcDNA3.1.

Immunoblotting and gene expression

Protein isolation and immunoblotting procedures were as described previously [54]. For relative quantification of APP cleavage by alpha-secretase (sAPP α) versus beta-secretase (sAPP β) pathways, cells were incubated for between 20-24 hours in Optimem (Gibco), and media concentrated (using 30 kDa Amicon Ultra 15 ml filters; Merck Millipore, Livingston, UK) by centrifugation (4000 x g) and subjected to SDS-PAGE with amounts presented relative to total protein. Protein antibodies used were: anti-Actin (Sigma-Aldrich; 1:5000), anti-APP (Ab54, GSK; 1:4000), anti-sAPP α (Cambridge Bioscience, Cambridge, UK; 1:1000), anti-sAPP β (GSK; 1:1000), anti-BACE1 (Sigma-Aldrich; 1:1000), anti-BAD (New England Biolabs, Hitchin, UK; 1:1000), with anti-total PDH (1 mg/ml) and anti-pPDH ϵ 1 α (1 mg/ml) from Drug Discovery Unit, University of Dundee.

Glucose oxidation assay

SH-SY5Y_{EV}, SH-SY5Y_{B1} or SH-SY5Y_{mB1} cells were plated into 6-well cell culture plates and any pre-treatments carried out as described above and in the results section. To begin the assay, cells were washed twice with Hepes-buffered saline (HBS (in mM); 140 NaCl, 20 Hepes, 5 KCl, 2.5 MgSO₄ and 1 CaCl₂, pH 7.4) and incubated in HBS containing 2.5 mM glucose and 74 kBq/ml D-[U-¹⁴C]glucose (PerkinElmer) along with any relevant inhibitors or treatments indicated in the results section for 3 hours at 37 °C. Media were transferred and ¹⁴CO₂ liberated by acidification with 60% perchloric acid, trapped by Whatman (GF/B) filter papers discs pre-soaked with 1 M KOH and radioactivity quantified by liquid-scintillation counting. Cells from the assay were washed twice with ice-cold 0.9 % NaCl and lysed with 1ml of 50 mM NaOH, and the radioactivity contained within the lysate quantified by liquid-scintillation counting, which served as a measure of the ¹⁴C incorporation into the cell during the assay period. Total protein content was determined in the lysate via the Bradford method and used to normalize glucose incorporation and oxidation rates for each sample.

Cellular respiration

The Seahorse Extracellular Flux Analyser utilises solid sensors that simultaneously monitor the oxygenation and pH of the media. The rate of oxygen consumption (OCR) and extracellular acidification rates (ECAR) can therefore be assessed in near real time allowing for high resolution changes in a range of metabolic parameters. SH-SY5Y_{EV}, SH-SY5Y_{B1} or SH-SY5Y_{mB1} cell monolayers were seeded into XF 24-well culture microplates (Seahorse Bioscience, Copenhagen, Denmark) the day prior to treatment or assay as indicated in the

results section. Optimal cell number was determined following a cell titration assay taking into account oxygen consumption rate (OCR) and extracellular acidification rate (ECAR), oxygen tension values and the appearance of the cell monolayers and was determined to be 40,000 cells. On the day of the assay, cells were placed in unbuffered DMEM with relevant inhibitors/treatments as indicated and placed in a non-CO₂ incubator for 1 hour (to de-gas solutions) prior to assay initiation. Standard 3 minute mix, 2 minute wait and 3 minute measure cycles were used; with 5 baseline measurements taken before, and a subsequent 3 measurements acquired following, drug additions.

Enzyme activity assays

Activity assays for pyruvate dehydrogenase (PDH; Abcam, Cambridge, UK), α -ketoglutarate dehydrogenase (α -KGDH; Antibodies Online, Aachen, Germany), isocitrate dehydrogenase (IDH; Sigma-Aldrich) and fumarase (Abcam) were performed according to the manufacturer's instructions.

Results

Overexpression of BACE1 increases amyloidogenic APP processing and suppresses glucose oxidation

Control, empty vector-treated (SH-SY5Y_{EV}) and BACE1 overexpressing (SH-SY5Y_{B1}) cells displayed equivalent APP protein levels, showing the three predominant APP transcript protein isoforms present in neurons ([55]; Fig. 1A,B). As expected, SH-SY5Y_{B1} cells exhibited significantly higher BACE1 protein levels (Fig. 1A,C), which resulted in altered APP cleavage; promoting a substantial shift from non-amyloidogenic to amyloidogenic processing (as denoted by increased sAPP β and decreased sAPP α levels (Fig. 1A,D,E). Overexpression of BACE1 resulted in a marked reduction in the rate of ¹⁴C-glucose oxidation, compared to SH-SY5Y_{EV} cells (Fig. 1F), in agreement with recent findings in C₂C₁₂ skeletal muscle cells [34]. The decrease in glucose oxidation is not due to impaired glucose incorporation into the cell (Fig. 1G), and suggests that increased BACE1 protein levels and activity cause a fundamental change in the ability of SH-SY5Y cells to oxidise glucose (Fig. 1H).

In an attempt to discern whether the changes in glucose oxidation rate were dependent upon the secretase activity of BACE1, a concomitant group of cells that overexpress a mutant form of BACE1, devoid of secretase activity (SH-SY5Y_{mB1}; [56]) were examined. BACE1 protein levels in SH-SY5Y_{mB1} cells were raised to a similar extent to that observed for SH-SY5Y_{B1} cells, with no alteration in APP protein levels (Fig. 2A-C). Interestingly, the overexpression of the secretase-dead BACE1 mutant exerted a dominant negative effect over APP processing with reduced sAPP β release, compared with SH-SY5Y_{B1} and SH-SY5Y_{mB1} cells, into the culture media (Fig. 2A,D). The reduced β -secretase cleavage of APP in the presence of mBACE1 was also mirrored in its effect on glucose oxidation rate, with no reduction in glucose oxidation concomitant with comparable glucose incorporation between cell types (Fig. 2E,F), indicating that increased BACE1 activity is required to depress glucose oxidation in SH-SY5Y cells (Fig. 2G).

Chronic elevation of BACE1 stimulates aerobic glycolysis in SH-SY5Y cells

The cellular pathways that facilitate glucose metabolism are glycolysis (and the pentose phosphate pathway), which occurs in the cytoplasm and the TCA cycle and oxidative phosphorylation, which are present in the matrix and inner mitochondrial membrane, respectively. Actively respiring mitochondria consume oxygen and therefore oxygen consumption rate (OCR) can be taken as a measure of substrate flux through the oxidative phosphorylation pathway while extracellular acidification rate (ECAR) reflects the release of lactate (lactic acid) converted from pyruvate following glycolysis. SH-SY5Y_{B1} cells exhibit decreased OCR concurrent with increased ECAR in comparison to SH-SY5Y_{EV} cells (Fig. 3A,B). Taken together these changes reflect a robust shift between these metabolic processes as shown by the change in the ratio of oxidative phosphorylation to glycolytic metabolism in cells with increased BACE1 activity (Fig. 3C).

To further investigate these changes, a glycolysis stress test (which assesses basal glycolysis, glycolytic capacity and glycolytic reserve) was performed (Fig. 3D). SH-SY5Y_{EV} and SH-SY5Y_{B1} cells were incubated in media containing zero glucose, 2.5 mM pyruvate and 4 mM L-glutamine for 100 minutes, following which stimulation of basal glycolysis was achieved by the injection of 2.5 mM glucose into the assay media. The increase in basal ECAR was significantly higher in SH-SY5Y_{B1} cells, compared to control cells, indicating that the raised BACE1 activity augmented aerobic glycolysis (Fig. 3E). Inhibition of F₁F₀ ATP synthase by oligomycin leaves cells wholly reliant on glycolysis for ATP generation (termed maximal glycolysis). This rate was also significantly increased in SH-SY5Y_{B1}, compared to control, cells following BACE1 overexpression (Fig. 3F). Finally, the cellular glycolytic reserve is given by the difference between basal and maximal glycolysis, with a significantly higher glycolytic reserve observed in the SH-SY5Y_{B1} cells (Fig. 3G). Collectively, these data indicate that raised BACE1 activity depresses glucose oxidation in SH-SY5Y cells and, as a result, there is a compensatory increase in glucose metabolism through aerobic glycolysis.

Mitochondrial efficiency is unaltered by raised BACE1 activity in SH-SY5Y cells

The marked reduction in oxidative phosphorylation, concurrent with increased glycolytic metabolism in SH-SY5Y_{B1} cells, indicated that increased BACE1 activity reduced substrate delivery to the mitochondria and/or impaired mitochondrial function. To test mitochondrial efficiency in these cells, a modified Mitochondrial Stress Test (Seahorse Bioscience) protocol was utilised. As previously reported, a limited maximal respiration rate is achievable in undifferentiated SH-SY5Y cells [57] in the absence of pyruvate in the assay media. Consequently, we therefore performed a split assay to investigate the effects of mitochondrial metabolism perturbation: the first part measuring the proportion of respiration dedicated to ATP generation (ATP synthase inhibition with oligomycin), the maintenance of mitochondrial leak (a combination of rotenone (complex I inhibition) and antimycin A (complex III inhibition) to minimize mitochondrial respiration) and the relative contribution of non-mitochondrial OCR (Fig. 4A). However, increased BACE1 activity in SH-SY5Y cells had no effect on any of these oxidative parameters (Fig. 4B-D). The second part encompasses a test of the cellular reserve capacity via the induction of maximal respiration through addition of the proton ionophore FCCP. This drug collapses the mitochondrial membrane potential, leading to uncoupled (no ATP generation) substrate flux through the mitochondria, promoting an increase in glycolysis in an attempt to maintain intracellular ATP levels. Raised BACE1 activity also had no effect on SH-SY5Y mitochondrial reserve capacity (Fig. 4E,F). The results from these assays show that chronic elevation of BACE1 protein expression and activity does not impair mitochondrial electron transfer function.

Consequently, this result strongly indicated that the BACE1 activity-driven decrease in glucose oxidation was the result of diminished substrate delivery to the mitochondria.

Raised BACE1 activity lesions key metabolic pathways in oxidative glucose metabolism

There are three key control points involved in the regulation of neuronal metabolism, these are the generation of glucose-6-phosphate, pyruvate and acetyl CoA. In neurons, pyruvate is predominantly produced directly from metabolism of glucose or indirectly, through the provision of extracellular lactate via the astrocyte-neuron lactate shuttle [58,59]. Therefore, in an attempt to better understand the changes in cellular metabolism induced by increased BACE1 activity, lactate utilisation by SH-SY5Y cells was assessed. SH-SY5Y_{B1} cells displayed significantly reduced lactate consumption as assessed by the ability of lactate to repress ¹⁴C-labelled glucose oxidation (Fig. 5A). This deficit could also be observed as a reduction in OCR when cells were provided physiological concentrations (0.5 or 2 mM; [60]) of lactate as sole substrate, although the reduction in lactate metabolism was overcome by the presentation of a higher (4 mM) lactate concentration (Fig. 5A,B). As the oxidation of both glucose and lactate was diminished, we hypothesised that raised BACE1 activity resulted in impairment of the pyruvate dehydrogenase complex activity (PDH). To address this we used an indirect PDH activity assay and demonstrated that SH-SY5Y_{B1} cells displayed a marked reduction of PDH activity (Fig. 5C). Consistent with this outcome, SH-SY5Y_{B1} cells displayed reduced OCR when provided with pyruvate (2.5 mM) as the sole substrate (Fig. 5D).

In response to cellular injury or stress, compensatory metabolic strategies are employed in an attempt to bypass reduced glucose utilization associated with decreased PDH activity and obviate impaired mitochondrial TCA function. For example, glutamine metabolism is diverted from an oxidative to reductive route [61,62], causing increased conversion to glutamate and, through glutamate dehydrogenase, raised levels of α -ketoglutarate. This additional source of α -ketoglutarate should supplement ATP generation through the TCA cycle as it bypasses the block at PDH. Consequently, we examined whether such an alternative route for TCA substrate replenishment was capable of recovering oxidative metabolism. Thus OCR and ECAR were measured when SH-SY5Y_{EV} and SH-SY5Y_{B1} cells were provided glucose (2.5 mM) alone or glucose (2.5 mM) + glutamine (4 mM). However, the presence of glutamine did not affect the attenuation of OCR or enhancement of ECAR in SH-SY5Y_{B1} cells (Fig. 5E). Moreover, OCR was reduced in SH-SY5Y_{B1} cells when glutamine was provided as the exclusive substrate (Fig. 5F). These data indicate that either glutamine does not play a significant role in SH-SY5Y basal metabolism and/or that raised BACE1 activity blunts additional TCA enzymes, such as α -ketoglutarate dehydrogenase (α -KGDH), which converts α -ketoglutarate to succinyl-CoA. Indeed, SH-SY5Y_{B1} cells showed a large, BACE1 secretase activity-dependent, reduction in α -KGDH activity, which is the rate-controlling step of the TCA cycle (Fig. 5G).

Because PDH and α -KGDH catalyse decarboxylation reactions involved in mitochondrial bioenergetics, we assayed the activity of the third enzyme utilising this process, isocitrate dehydrogenase (IDH). Three isoforms of IDH are present in cells, IDH3 in the mitochondrial matrix, which reduces NAD⁺ to NADH, and IDH1 (cytoplasmic and peroxisomal) and IDH2 (mitochondrial) reducing NADP⁺ to NADPH. Consequently, specific isoform function was determined by differential supplementation (NAD⁺ or NADP⁺) of the reaction mixes. In agreement with the results for the decarboxylation enzymes above, raised BACE1 activity

resulted in depression of NAD⁺- and NADP⁺-dependent IHD activity (Fig. 5H). However impairment of TCA cycle enzyme activities by increased BACE1 activity was not universal. For example, although fumarase activity was reduced slightly in SH-SY5Y_{B1} cells, this was matched in SH-SY5Y_{mB1} cells, with the protein levels of fumarase also decreased in both these cell lines, resulting in unaltered specific activity of fumarase by raised BACE1 in SH-SY5Y cells (Fig. 5I). Thus manipulation of APP cleavage towards amyloidogenic processing by increased BACE1 activity in SH-SY5Y cells reduces the overall catabolic capacity of the TCA cycle through specific enzyme lesions rather than a wholesale down-regulation.

Rescue of glucose oxidation in SH-SY5Y_{B1} cells.

As glutamine supplementation was unable to recover BACE1-mediated inhibition of glucose oxidation, we focused on PDH as a potential target for pharmacological or alternative nutrient interventions. A key regulator of PDH activity is by phosphorylation of its E1 α subunit [63], which is predominantly controlled by the inhibitory effect of pyruvate dehydrogenase kinases (PDKs) versus activation through de-phosphorylation by pyruvate dehydrogenase phosphatase (PDP). Dichloroacetate (DCA) is a PDK1 inhibitor, which enhances oxidative phosphorylation in the brain [58] and has been promoted clinically for its anti-neoplastic effects [64]. Consistent with PDH inhibition, SH-SY5Y_{B1} cells exhibited increased phosphorylation of the e1 α subunit at serine 293 compared to SH-SY5Y_{EV} cells (Fig. 6A). Treatment with DCA (10 and 100 μ M) resulted in decreased levels of Ser²⁹³ e1 α phosphorylation of both SH-SY5Y cell types, although Ser²⁹³ e1 α phosphorylation remained significantly higher in SH-SY5Y_{B1} cells (Fig. 6A,B). In agreement with the reduction in e1 α phosphorylation, incubation of SH-SY5Y_{EV} cells with DCA (100 μ M) increased the glucose oxidation rate (Fig. 6C). However, although glucose oxidation of SH-SY5Y_{B1} cells was also enhanced by DCA treatment, this remained significantly lower than that of the SH-SY5Y_{EV} cells, mirroring e1 α phosphorylation status.

An alternative means of bypassing PDH is through the application of ketones, such as beta-hydroxybutyrate (BHB), which is metabolised to acetyl-CoA and enters the TCA cycle at oxaloacetate. The presence of BHB (0.5 – 10 μ M) recovered the relative deficit in glucose oxidation of SH-SY5Y_{B1} cells, compared to SH-SY5Y_{EV} cells, in a concentration-dependent manner (Fig. 6D). Previous studies [65,66] have indicated that diminished mitochondrial consumption of glucose in neurons and the ability to switch substrate preference to ketones is dependent on the presence and activity of BAD (BCL-2-associated agonist of cell death), a member of the BCL-2 gene family member of apoptotic control proteins, with increased mitochondrial usage of BHB in *Bad*^{-/-} cortical neurons. In agreement with this model, we find that the increased ability of SH-SY5Y_{B1} cells to utilise BHB over SH-SY5Y_{EV} cells and enhance glucose oxidation, is associated with a large reduction of BAD protein expression (Fig. 6E). Finally we also investigated the ability of the naturally occurring enzyme co-factor, α -lipoic acid to modify glucose oxidation of SH-SY5Y_{B1} cells as previous studies have shown this compound to up-regulate mitochondrial bioenergetics and promote glucose uptake [67,68]. Indeed PDH and α -KGDH protein complexes require α -lipoic acid as a co-factor for their acyl transferase activity. Supplementation of the growth media with α -lipoic acid (25 or 50 μ M for 48 hours) resulted in a robust concentration-dependent attenuation of the impaired glucose oxidation rate displayed by SH-SY5Y_{B1}, compared to SH-SY5Y_{EV} cells (Fig. 6F). Taken together, these results demonstrate the applicability of alternative nutrient (BHB) or co-factor (α -lipoic acid) supplementation to alleviate or by-pass the impaired glucose metabolism elicited by raised BACE1 activity in SH-SY5Y cells.

358

359 Discussion

360 In this study we demonstrate that raised BACE1 activity by overexpression in SH-SY5Y
361 cells, and subsequent modification of APP metabolism, induces a shift from the
362 physiologically predominant alpha-secretase cleavage pathway to the amyloidogenic beta-
363 secretase cleavage pathway, which results in decreased glucose metabolism. This
364 bioenergetic impairment was dependent upon the secretase activity of BACE1 and produced
365 a fundamental shift in the cellular metabolic profile, characterised by reduced glucose
366 oxidation in association with a compensatory increase in glycolysis, in an attempt to maintain
367 ATP production.

368 Decreased brain glucose metabolism is an invariant pathophysiological event occurring in
369 AD progression, and it has been hypothesised that this may occur years, and even decades
370 prior to symptom presentation [15,16,69,70]. Impaired glucose metabolism has also been
371 noted in central tissues taken from AD animal models [22,23,68]. Furthermore, despite the
372 observation that reduced glucose metabolism is predictive of later cognitive decline and AD
373 symptom presentation, relatively little is known about the cellular mechanisms underlying
374 these changes. Interestingly, the metabolic shift we have observed in favour of aerobic
375 glycolysis under conditions of increased BACE1 activity mirrors early and predictive
376 changes occurring in the brains of people who later develop AD. The brain areas that display
377 this profile overlap with regions showing the greatest prevalence of A β pathology, future
378 susceptibility to cell death and are predictive of cognitive decline [32,33].

379 The mechanisms underlying glucose hypometabolism in AD are not well understood.
380 However mitochondrial dysfunction has been widely reported in clinical and experimental
381 AD studies [19-21] and A β has been reported to accumulate in the mitochondria of AD
382 patients and transgenic AD mouse models prior to amyloid deposition [25-27,71]. Although
383 there are numerous reports of impaired electron transfer in mitochondria, for example via
384 diminished activity in complexes I to V in AD subjects [72-75], and diminished complex III
385 and IV activity in a transgenic AD mouse model [76] we found no effect of raised BACE1
386 activity on mitochondrial electron transfer function in SH-SY5Y cells. It may be that
387 additional injurious processes, such as increased oxidative and/or inflammatory stress are
388 required to act concurrently with raised BACE1 activity and increased levels of A β to elicit
389 this outcome.

390 Thus the driving force for decreased glucose oxidation in SH-SY5Y_{B1} cells appears to be
391 through reduced substrate delivery to mitochondria by a BACE1 activity-dependent
392 impairment of specific TCA cycle enzymes; PDH, α KGDH and IDH each of which utilise
393 decarboxylation reactions. Previous studies on post-mortem AD brains indicate reduced
394 levels and/or activity of these key enzymes: PDH, α KGDH and IDH [20,77-81], with PDH
395 and α KGDH exhibiting the largest decreases in activity. Furthermore, A β peptides have been
396 demonstrated to inhibit PDH and α KGDH directly [28,31,82]. In contrast we find no change
397 in the activity of the TCA enzyme fumarase, the activity of which is also unaltered in AD
398 brains [20]. Our results add to these findings by implicating raised BACE1 activity and
399 manipulation of APP processing as central to these enzyme deficits and metabolic adaptations
400 at the cellular level, indicating a key role for BACE1 and its up-regulation in response to
401 oxidative and inflammatory stress, which are associated with the very early stages of AD
402 [39,43,83]. Indeed, these results show that raised BACE1 activity effectively phenocopies

some of the earliest changes in glucose metabolism seen in the brain during the progression towards AD.

In situations of diminished glucose utilization, there are a number of strategies that may be engaged in an attempt to bypass this block in metabolism. For example, in brain ischemic-reperfusion injury there is also decline in glucose oxidation and PDH activity [84,85]. Application of glutamine has been demonstrated to offer some protection against ischemic-reperfusion injury in peripheral tissues by providing an alternative energy source for the cell to maintain ATP content [86,87]. In addition to providing a source of carbon for neurotransmitter production, glutamine can also feed into the TCA cycle and be metabolised to increase levels of α -ketoglutarate and succinate. However, our data demonstrate that glutamine supplementation is unable to restore oxidative glucose metabolism in cells overexpressing BACE1. Furthermore, SH-SY5Y_{B1} cells exhibited diminished oxidation of glutamine when applied as sole substrate, in line with the reduced activity of α -KGDH observed in these cells.

The inhibition of PDH activity in SH-SY5Y_{B1} cells was associated with increased phosphorylation of the e1 α subunit at Ser²⁹³, which is the main inhibitory site for PDH activity [88]. Although PDK1 inhibition with dichloroacetate reduced e1 α phosphorylation levels, this was also observed in SH-SY5Y_{EV} cells, in conjunction with a maintained relative lower glucose oxidation in SH-SY5Y_{B1} cells. This outcome may be owing to the presence of BACE1 activity-sensitive DCA-resistant PDK isoenzyme, or PDP, or an alternative kinase that phosphorylates e1 α at Ser²⁹³ (e.g. GSK3). In contrast, application of the ketone, BHB, to SH-SY5Y_{B1} cells demonstrates complete recovery of glucose oxidation, relative to SH-SY5Y_{EV} cells, presumably by directly increasing the availability of acetyl CoA to the TCA cycle, thus circumventing PDH. The enhanced ability to utilise ketone bodies to aid glucose oxidation in SH-SY5Y_{B1} cells may result from their reduced BAD levels, compared to SH-SY5Y_{EV} cells. This finding agrees with the work of Nikita Danial's group showing that neurons derived from BAD knock out animals display an augmented bioenergetic profile in the presence of ketone bodies [66].

α -lipoic acid is synthesised in mitochondria and is a necessary cofactor for PDH and α -KGDH activity. Our finding that exogenously applied α -lipoic acid also increases glucose oxidation in SH-SY5Y_{B1}, but not SH-SY5Y_{EV} cells suggests that raised BACE1 activity either reduces mitochondrial α -lipoic acid levels or lessens the ability of α -lipoic acid to enhance PDH and α -KGDH activity. The latter mechanism may be favoured as there is evidence that excess α -lipoic acid provides protection to neuronal cells from the deleterious effects of exogenously applied A β peptides on PDH activity [89] and A β -induced toxicity [90]. Regarding the potential to reverse the metabolic deficits associated with age-dependent cognitive decline and early-stage or mild AD, the data presented herein demonstrate the capacity of α -lipoic acid and BHB to attenuate impaired glucose oxidation at a cellular level in face of chronically raised BACE1 activity and increased levels of A β peptides. BHB and α -lipoic acid have been demonstrated to exhibit some positive effects on cognitive decline in elderly dementia and mild AD patients [11,13,91-93].

Taken together, our findings are supportive for the rationale of targeting these TCA enzyme deficits in early cognitive impairment and AD. However, so far clinical trials based on raising circulating levels of ketones or giving α -lipoic acid as a dietary supplement have provided mixed outcomes and further trials examining α -lipoic acid are currently in progress. Perhaps a future strategy for AD management and treatment may be through the combination

of this nutraceutical approach with new therapeutics, such as BACE1 inhibitors, currently in clinical trials [94].

Acknowledgements

The authors thank Dundee Cell Products and Professor Wolfe for help with the cloning and provision of the mBACE1 construct, respectively. This work was supported by the Cruden Foundation Ltd and by grants from Alzheimer's Research UK (ART-PhD2010-12 and ARUK-EG2012A-2) and Diabetes UK (0003681).

Author contributions

JAF and DLH performed experiments and analysed data. JAF, DLH and MLJA contributed to the conception and design of experiments, interpretation of data and drafting and revising the manuscript. MLJA supervised the study and JAF and MLJA wrote the manuscript. All authors approved the final version. MLJA is the guarantor of this work

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Jack CR, Jr., Knopman DS, Jagust WJ, Petersen RC, Weiner MW, Aisen PS, Shaw LM, Vemuri P, Wiste HJ, Weigand SD, Lesnick TG, Pankratz VS, Donohue MC, Trojanowski JQ (2013) Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol* 12 (2):207-216. doi:10.1016/S1474-4422(12)70291-0
2. Jack CR, Jr., Wiste HJ, Weigand SD, Knopman DS, Lowe V, Vemuri P, Mielke MM, Jones DT, Senjem ML, Gunter JL, Gregg BE, Pankratz VS, Petersen RC (2013) Amyloid-first and neurodegeneration-first profiles characterize incident amyloid PET positivity. *Neurology* 81 (20):1732-1740. doi:10.1212/01.wnl.0000435556.21319.e4
3. Knopman DS, Jack CR, Jr., Wiste HJ, Weigand SD, Vemuri P, Lowe VJ, Kantarci K, Gunter JL, Senjem ML, Mielke MM, Roberts RO, Boeve BF, Petersen RC (2013) Selective worsening of brain injury biomarker abnormalities in cognitively normal elderly persons with beta-amyloidosis. *JAMA Neurol* 70 (8):1030-1038. doi:10.1001/jamaneurol.2013.182
4. Martiskainen H, Helisalmi S, Viswanathan J, Kurki M, Hall A, Herukka SK, Sarajarvi T, Natunen T, Kurkinen KM, Huovinen J, Makinen P, Laitinen M, Koivisto AM, Mattila KM, Lehtimäki T, Remes AM, Leinonen V, Haapasalo A, Soininen H, Hiltunen M (2015) Effects of Alzheimer's disease-associated risk loci on cerebrospinal fluid biomarkers and disease progression: a polygenic risk score approach. *J Alzheimers Dis* 43 (2):565-573. doi:10.3233/JAD-140777
5. Swerdlow RH, Burns JM, Khan SM (2014) The Alzheimer's disease mitochondrial cascade hypothesis: progress and perspectives. *Biochim Biophys Acta* 1842 (8):1219-1231. doi:10.1016/j.bbadis.2013.09.010
6. Wilkins HM, Harris JL, Carl SM, E L, Lu J, Eva Selfridge J, Roy N, Hutfles L, Koppel S, Morris J, Burns JM, Michaelis ML, Michaelis EK, Brooks WM, Swerdlow RH (2014) Oxaloacetate activates brain mitochondrial biogenesis, enhances the insulin pathway, reduces inflammation and stimulates neurogenesis. *Hum Mol Genet* 23 (24):6528-6541. doi:10.1093/hmg/ddu371
7. Yao J, Irwin RW, Zhao L, Nilsen J, Hamilton RT, Brinton RD (2009) Mitochondrial bioenergetic deficit precedes Alzheimer's pathology in female mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 106 (34):14670-14675. doi:10.1073/pnas.0903563106
8. Bouzier-Sore AK, Voisin P, Bouchaud V, Bezancon E, Franconi JM, Pellerin L (2006) Competition between glucose and lactate as oxidative energy substrates in both neurons and astrocytes: a comparative NMR study. *Eur J Neurosci* 24 (6):1687-1694. doi:10.1111/j.1460-9568.2006.05056.x

9. McKenna MC, Waagepetersen HS, Schousboe A, Sonnewald U (2006) Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: current evidence and pharmacological tools. *Biochem Pharmacol* 71 (4):399-407. doi:10.1016/j.bcp.2005.10.011
10. Sokoloff L (1999) Energetics of functional activation in neural tissues. *Neurochem Res* 24 (2):321-329
11. Henderson ST, Vogel JL, Barr LJ, Garvin F, Jones JJ, Costantini LC (2009) Study of the ketogenic agent AC-1202 in mild to moderate Alzheimer's disease: a randomized, double-blind, placebo-controlled, multicenter trial. *Nutr Metab (Lond)* 6:31. doi:10.1186/1743-7075-6-31
12. Blass JP (2001) Brain metabolism and brain disease: is metabolic deficiency the proximate cause of Alzheimer dementia? *J Neurosci Res* 66 (5):851-856
13. Costantini LC, Barr LJ, Vogel JL, Henderson ST (2008) Hypometabolism as a therapeutic target in Alzheimer's disease. *BMC Neurosci* 9 Suppl 2:S16. doi:10.1186/1471-2202-9-S2-S16
14. Johnson KA, Lopera F, Jones K, Becker A, Sperling R, Hilson J, Londono J, Siegert I, Arcos M, Moreno S, Madrigal L, Ossa J, Pineda N, Ardila A, Roselli M, Albert MS, Kosik KS, Rios A (2001) Presenilin-1-associated abnormalities in regional cerebral perfusion. *Neurology* 56 (11):1545-1551
15. Mosconi L, Mistur R, Switalski R, Brys M, Glodzik L, Rich K, Pirraglia E, Tsui W, De Santi S, de Leon MJ (2009) Declining brain glucose metabolism in normal individuals with a maternal history of Alzheimer disease. *Neurology* 72 (6):513-520. doi:10.1212/01.wnl.0000333247.51383.43
16. Mosconi L, Mistur R, Switalski R, Tsui WH, Glodzik L, Li Y, Pirraglia E, De Santi S, Reisberg B, Wisniewski T, de Leon MJ (2009) FDG-PET changes in brain glucose metabolism from normal cognition to pathologically verified Alzheimer's disease. *Eur J Nucl Med Mol Imaging* 36 (5):811-822. doi:10.1007/s00259-008-1039-z
17. Reiman EM, Chen K, Alexander GE, Caselli RJ, Bandy D, Osborne D, Saunders AM, Hardy J (2004) Functional brain abnormalities in young adults at genetic risk for late-onset Alzheimer's dementia. *Proc Natl Acad Sci U S A* 101 (1):284-289. doi:10.1073/pnas.2635903100
18. Small GW, Mazziotta JC, Collins MT, Baxter LR, Phelps ME, Mandelkern MA, Kaplan A, La Rue A, Adamson CF, Chang L, et al. (1995) Apolipoprotein E type 4 allele and cerebral glucose metabolism in relatives at risk for familial Alzheimer disease. *JAMA* 273 (12):942-947
19. Beal MF (2005) Oxidative damage as an early marker of Alzheimer's disease and mild cognitive impairment. *Neurobiol Aging* 26 (5):585-586. doi:10.1016/j.neurobiolaging.2004.09.022
20. Bubber P, Haroutunian V, Fisch G, Blass JP, Gibson GE (2005) Mitochondrial abnormalities in Alzheimer brain: mechanistic implications. *Ann Neurol* 57 (5):695-703. doi:10.1002/ana.20474
21. Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu S, Gunn-Moore F, Lue LF, Walker DG, Kuppusamy P, Zewier ZL, Arancio O, Stern D, Yan SS, Wu H (2004) AβAD directly links Aβeta to mitochondrial toxicity in Alzheimer's disease. *Science* 304 (5669):448-452. doi:10.1126/science.1091230
22. Nilsen LH, Witter MP, Sonnewald U (2014) Neuronal and astrocytic metabolism in a transgenic rat model of Alzheimer's disease. *J Cereb Blood Flow Metab* 34 (5):906-914. doi:10.1038/jcbfm.2014.37
23. von Kienlin M, Kunnecke B, Metzger F, Steiner G, Richards JG, Ozmen L, Jacobsen H, Loetscher H (2005) Altered metabolic profile in the frontal cortex of PS2APP transgenic mice, monitored throughout their life span. *Neurobiol Dis* 18 (1):32-39. doi:10.1016/j.nbd.2004.09.005
24. Yao J, Chen S, Mao Z, Cadenas E, Brinton RD (2011) 2-Deoxy-D-glucose treatment induces ketogenesis, sustains mitochondrial function, and reduces pathology in female mouse model of Alzheimer's disease. *PLoS One* 6 (7):e21788. doi:10.1371/journal.pone.0021788
25. Devi L, Prabhu BM, Galati DF, Avadhani NG, Anandatheerthavarada HK (2006) Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. *J Neurosci* 26 (35):9057-9068. doi:10.1523/JNEUROSCI.1469-06.2006

549 26. Hirai K, Aliev G, Nunomura A, Fujioka H, Russell RL, Atwood CS, Johnson AB, Kress Y, Vinters HV,
550 Tabaton M, Shimohama S, Cash AD, Siedlak SL, Harris PL, Jones PK, Petersen RB, Perry G, Smith MA
551 (2001) Mitochondrial abnormalities in Alzheimer's disease. *J Neurosci* 21 (9):3017-3023
552 27. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH (2006) Mitochondria are a
553 direct site of A beta accumulation in Alzheimer's disease neurons: implications for free radical
554 generation and oxidative damage in disease progression. *Hum Mol Genet* 15 (9):1437-1449.
555 doi:10.1093/hmg/ddl066
556 28. Casley CS, Canevari L, Land JM, Clark JB, Sharpe MA (2002) Beta-amyloid inhibits integrated
557 mitochondrial respiration and key enzyme activities. *J Neurochem* 80 (1):91-100
558 29. Caspersen C, Wang N, Yao J, Sosunov A, Chen X, Lustbader JW, Xu HW, Stern D, McKhann G, Yan
559 SD (2005) Mitochondrial Abeta: a potential focal point for neuronal metabolic dysfunction in
560 Alzheimer's disease. *FASEB J* 19 (14):2040-2041. doi:10.1096/fj.05-3735fje
561 30. Pereira C, Ferreira E, Cardoso SM, de Oliveira CR (2004) Cell degeneration induced by amyloid-
562 beta peptides: implications for Alzheimer's disease. *J Mol Neurosci* 23 (1-2):97-104.
563 doi:10.1385/JMN:23:1-2:097
564 31. Shearman MS, Ragan CI, Iversen LL (1994) Inhibition of PC12 cell redox activity is a specific, early
565 indicator of the mechanism of beta-amyloid-mediated cell death. *Proc Natl Acad Sci U S A* 91
566 (4):1470-1474
567 32. Vaishnavi SN, Vlassenko AG, Rundle MM, Snyder AZ, Mintun MA, Raichle ME (2010) Regional
568 aerobic glycolysis in the human brain. *Proc Natl Acad Sci U S A* 107 (41):17757-17762.
569 doi:10.1073/pnas.1010459107
570 33. Vlassenko AG, Vaishnavi SN, Couture L, Sacco D, Shannon BJ, Mach RH, Morris JC, Raichle ME,
571 Mintun MA (2010) Spatial correlation between brain aerobic glycolysis and amyloid-beta (Abeta)
572 deposition. *Proc Natl Acad Sci U S A* 107 (41):17763-17767. doi:10.1073/pnas.1010461107
573 34. Hussain I, Powell D, Howlett DR, Tew DG, Meek TD, Chapman C, Gloger IS, Murphy KE, Southan
574 CD, Ryan DM, Smith TS, Simmons DL, Walsh FS, Dingwall C, Christie G (1999) Identification of a novel
575 aspartic protease (Asp 2) as beta-secretase. *Mol Cell Neurosci* 14 (6):419-427.
576 doi:10.1006/mcne.1999.0811
577 35. Sinha S, Anderson JP, Barbour R, Basi GS, Caccavello R, Davis D, Doan M, Dovey HF, Frigon N,
578 Hong J, Jacobson-Croak K, Jewett N, Keim P, Knops J, Lieberburg I, Power M, Tan H, Tatsuno G, Tung
579 J, Schenk D, Seubert P, Suomensaaari SM, Wang S, Walker D, Zhao J, McConlogue L, John V (1999)
580 Purification and cloning of amyloid precursor protein beta-secretase from human brain. *Nature* 402
581 (6761):537-540. doi:10.1038/990114
582 36. Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, Denis P, Teplow DB, Ross S, Amarante P,
583 Loeloff R, Luo Y, Fisher S, Fuller J, Edenson S, Lile J, Jarosinski MA, Biere AL, Curran E, Burgess T, Louis
584 JC, Collins F, Treanor J, Rogers G, Citron M (1999) Beta-secretase cleavage of Alzheimer's amyloid
585 precursor protein by the transmembrane aspartic protease BACE. *Science* 286 (5440):735-741
586 37. Meakin PJ, Harper AJ, Hamilton DL, Gallagher J, McNeilly AD, Burgess LA, Vaanholt LM, Bannon
587 KA, Latcham J, Hussain I, Speakman JR, Howlett DR, Ashford ML (2012) Reduction in BACE1
588 decreases body weight, protects against diet-induced obesity and enhances insulin sensitivity in
589 mice. *Biochem J* 441 (1):285-296. doi:10.1042/BJ20110512
590 38. Hamilton DL, Findlay JA, Montagut G, Meakin PJ, Bestow D, Jalicy SM, Ashford ML (2014) Altered
591 amyloid precursor protein processing regulates glucose uptake and oxidation in cultured rodent
592 myotubes. *Diabetologia* 57 (8):1684-1692. doi:10.1007/s00125-014-3269-x
593 39. Guglielmotto M, Aragno M, Autelli R, Giliberto L, Novo E, Colombatto S, Danni O, Parola M, Smith
594 MA, Perry G, Tamagno E, Tabaton M (2009) The up-regulation of BACE1 mediated by hypoxia and
595 ischemic injury: role of oxidative stress and HIF1alpha. *J Neurochem* 108 (4):1045-1056.
596 doi:10.1111/j.1471-4159.2008.05858.x
597 40. O'Connor T, Sadleir KR, Maus E, Velliquette RA, Zhao J, Cole SL, Eimer WA, Hitt B, Bembinster LA,
598 Lammich S, Lichtenthaler SF, Hebert SS, De Strooper B, Haass C, Bennett DA, Vassar R (2008)

599 Phosphorylation of the translation initiation factor eIF2 α increases BACE1 levels and promotes
600 amyloidogenesis. *Neuron* 60 (6):988-1009. doi:10.1016/j.neuron.2008.10.047

601 41. Puglielli L, Ellis BC, Saunders AJ, Kovacs DM (2003) Ceramide stabilizes beta-site amyloid
602 precursor protein-cleaving enzyme 1 and promotes amyloid beta-peptide biogenesis. *J Biol Chem*
603 278 (22):19777-19783. doi:10.1074/jbc.M300466200

604 42. Velliquette RA, O'Connor T, Vassar R (2005) Energy inhibition elevates beta-secretase levels and
605 activity and is potentially amyloidogenic in APP transgenic mice: possible early events in Alzheimer's
606 disease pathogenesis. *J Neurosci* 25 (47):10874-10883. doi:10.1523/JNEUROSCI.2350-05.2005

607 43. Zhang X, Zhou K, Wang R, Cui J, Lipton SA, Liao FF, Xu H, Zhang YW (2007) Hypoxia-inducible
608 factor 1 α (HIF-1 α)-mediated hypoxia increases BACE1 expression and beta-amyloid
609 generation. *J Biol Chem* 282 (15):10873-10880. doi:10.1074/jbc.M608856200

610 44. Fang M, Wang J, Zhang X, Geng Y, Hu Z, Rudd JA, Ling S, Chen W, Han S (2012) The miR-124
611 regulates the expression of BACE1/beta-secretase correlated with cell death in Alzheimer's disease.
612 *Toxicol Lett* 209 (1):94-105. doi:10.1016/j.toxlet.2011.11.032

613 45. Lei X, Lei L, Zhang Z, Zhang Z, Cheng Y (2015) Downregulated miR-29c correlates with increased
614 BACE1 expression in sporadic Alzheimer's disease. *Int J Clin Exp Pathol* 8 (2):1565-1574

615 46. Nelson PT, Wang WX (2010) MiR-107 is reduced in Alzheimer's disease brain neocortex:
616 validation study. *J Alzheimers Dis* 21 (1):75-79. doi:10.3233/JAD-2010-091603

617 47. Fukumoto H, Cheung BS, Hyman BT, Irizarry MC (2002) Beta-secretase protein and activity are
618 increased in the neocortex in Alzheimer disease. *Arch Neurol* 59 (9):1381-1389

619 48. Holsinger RM, McLean CA, Beyreuther K, Masters CL, Evin G (2002) Increased expression of the
620 amyloid precursor beta-secretase in Alzheimer's disease. *Ann Neurol* 51 (6):783-786.
621 doi:10.1002/ana.10208

622 49. Tyler SJ, Dawbarn D, Wilcock GK, Allen SJ (2002) alpha- and beta-secretase: profound changes in
623 Alzheimer's disease. *Biochem Biophys Res Commun* 299 (3):373-376

624 50. Zhao J, Fu Y, Yasvoina M, Shao P, Hitt B, O'Connor T, Logan S, Maus E, Citron M, Berry R, Binder L,
625 Vassar R (2007) Beta-site amyloid precursor protein cleaving enzyme 1 levels become elevated in
626 neurons around amyloid plaques: implications for Alzheimer's disease pathogenesis. *J Neurosci* 27
627 (14):3639-3649. doi:10.1523/JNEUROSCI.4396-06.2007

628 51. Harada H, Tamaoka A, Ishii K, Shoji S, Kametaka S, Kametani F, Saito Y, Murayama S (2006) Beta-
629 site APP cleaving enzyme 1 (BACE1) is increased in remaining neurons in Alzheimer's disease brains.
630 *Neurosci Res* 54 (1):24-29. doi:10.1016/j.neures.2005.10.001

631 52. Li R, Lindholm K, Yang LB, Yue X, Citron M, Yan R, Beach T, Sue L, Sabbagh M, Cai H, Wong P,
632 Price D, Shen Y (2004) Amyloid beta peptide load is correlated with increased beta-secretase activity
633 in sporadic Alzheimer's disease patients. *Proc Natl Acad Sci U S A* 101 (10):3632-3637.
634 doi:10.1073/pnas.0205689101

635 53. Yang LB, Lindholm K, Yan R, Citron M, Xia W, Yang XL, Beach T, Sue L, Wong P, Price D, Li R, Shen
636 Y (2003) Elevated beta-secretase expression and enzymatic activity detected in sporadic Alzheimer
637 disease. *Nat Med* 9 (1):3-4. doi:10.1038/nm0103-3

638 54. Mirshamsi S, Laidlaw HA, Ning K, Anderson E, Burgess LA, Gray A, Sutherland C, Ashford ML
639 (2004) Leptin and insulin stimulation of signalling pathways in arcuate nucleus neurones: PI3K
640 dependent actin reorganization and KATP channel activation. *BMC Neurosci* 5:54. doi:10.1186/1471-
641 2202-5-54

642 55. Huang HC, Jiang ZF (2011) Amyloid-beta protein precursor family members: a review from
643 homology to biological function. *J Alzheimers Dis* 26 (4):607-626. doi:10.3233/JAD-2011-110335

644 56. Mowrer KR, Wolfe MS (2008) Promotion of BACE1 mRNA alternative splicing reduces amyloid
645 beta-peptide production. *J Biol Chem* 283 (27):18694-18701. doi:10.1074/jbc.M801322200

646 57. Xun Z, Lee DY, Lim J, Canaria CA, Barnebey A, Yanonne SM, McMurray CT (2012) Retinoic acid-
647 induced differentiation increases the rate of oxygen consumption and enhances the spare
648 respiratory capacity of mitochondria in SH-SY5Y cells. *Mech Ageing Dev* 133 (4):176-185.
649 doi:10.1016/j.mad.2012.01.008

58. Itoh Y, Esaki T, Shimoji K, Cook M, Law MJ, Kaufman E, Sokoloff L (2003) Dichloroacetate effects on glucose and lactate oxidation by neurons and astroglia in vitro and on glucose utilization by brain in vivo. *Proc Natl Acad Sci U S A* 100 (8):4879-4884. doi:10.1073/pnas.0831078100
59. Magistretti PJ, Pellerin L (1999) Cellular mechanisms of brain energy metabolism and their relevance to functional brain imaging. *Philos Trans R Soc Lond B Biol Sci* 354 (1387):1155-1163. doi:10.1098/rstb.1999.0471
60. Phipers B, Pierce JT (2006) Lactate physiology in health and disease. *Continuing Education in Anaesthesia, Critical Care & Pain* 6 (3):128-132. doi:10.1093/bjaceaccp/mkl018
61. Mullen AR, Wheaton WW, Jin ES, Chen PH, Sullivan LB, Cheng T, Yang Y, Linehan WM, Chandel NS, DeBerardinis RJ (2012) Reductive carboxylation supports growth in tumour cells with defective mitochondria. *Nature* 481 (7381):385-388. doi:10.1038/nature10642
62. Wise DR, Ward PS, Shay JE, Cross JR, Gruber JJ, Sachdeva UM, Platt JM, DeMatteo RG, Simon MC, Thompson CB (2011) Hypoxia promotes isocitrate dehydrogenase-dependent carboxylation of alpha-ketoglutarate to citrate to support cell growth and viability. *Proc Natl Acad Sci U S A* 108 (49):19611-19616. doi:10.1073/pnas.1117773108
63. Patel MS, Nemeria NS, Furey W, Jordan F (2014) The pyruvate dehydrogenase complexes: structure-based function and regulation. *J Biol Chem* 289 (24):16615-16623. doi:10.1074/jbc.R114.563148
64. Kankotia S, Stacpoole PW (2014) Dichloroacetate and cancer: new home for an orphan drug? *Biochim Biophys Acta* 1846 (2):617-629. doi:10.1016/j.bbcan.2014.08.005
65. Danial NN (2008) BAD: undertaker by night, candyman by day. *Oncogene* 27 Suppl 1:S53-70. doi:10.1038/onc.2009.44
66. Gimenez-Cassina A, Martinez-Francois JR, Fisher JK, Szlyk B, Polak K, Wiwczar J, Tanner GR, Lutas A, Yellen G, Danial NN (2012) BAD-dependent regulation of fuel metabolism and K(ATP) channel activity confers resistance to epileptic seizures. *Neuron* 74 (4):719-730. doi:10.1016/j.neuron.2012.03.032
67. Packer L, Cadenas E (2011) Lipoic acid: energy metabolism and redox regulation of transcription and cell signaling. *J Clin Biochem Nutr* 48 (1):26-32. doi:10.3164/jcbn.11-005FR
68. Sancheti H, Akopian G, Yin F, Brinton RD, Walsh JP, Cadenas E (2013) Age-dependent modulation of synaptic plasticity and insulin mimetic effect of lipoic acid on a mouse model of Alzheimer's disease. *PLoS One* 8 (7):e69830. doi:10.1371/journal.pone.0069830
69. Bateman RJ, Xiong C, Benzinger TL, Fagan AM, Goate A, Fox NC, Marcus DS, Cairns NJ, Xie X, Blazey TM, Holtzman DM, Santacruz A, Buckles V, Oliver A, Moulder K, Aisen PS, Ghetti B, Klunk WE, McDade E, Martins RN, Masters CL, Mayeux R, Ringman JM, Rossor MN, Schofield PR, Sperling RA, Salloway S, Morris JC, Dominantly Inherited Alzheimer N (2012) Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med* 367 (9):795-804. doi:10.1056/NEJMoa1202753
70. Jack CR, Jr., Barrio JR, Kepe V (2013) Cerebral amyloid PET imaging in Alzheimer's disease. *Acta Neuropathol* 126 (5):643-657. doi:10.1007/s00401-013-1185-7
71. Du H, Guo L, Fang F, Chen D, Sosunov AA, McKhann GM, Yan Y, Wang C, Zhang H, Molkentin JD, Gunn-Moore FJ, Vonsattel JP, Arancio O, Chen JX, Yan SD (2008) Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med* 14 (10):1097-1105. doi:10.1038/nm.1868
72. Bosetti F, Brizzi F, Barogi S, Mancuso M, Siciliano G, Tendi EA, Murri L, Rapoport SI, Solaini G (2002) Cytochrome c oxidase and mitochondrial F1F0-ATPase (ATP synthase) activities in platelets and brain from patients with Alzheimer's disease. *Neurobiol Aging* 23 (3):371-376
73. Kim SH, Vlkolinsky R, Cairns N, Fountoulakis M, Lubec G (2001) The reduction of NADH ubiquinone oxidoreductase 24- and 75-kDa subunits in brains of patients with Down syndrome and Alzheimer's disease. *Life Sci* 68 (24):2741-2750
74. Maurer I, Zierz S, Moller HJ (2000) A selective defect of cytochrome c oxidase is present in brain of Alzheimer disease patients. *Neurobiol Aging* 21 (3):455-462

75. Parker WD, Jr., Parks J, Filley CM, Kleinschmidt-DeMasters BK (1994) Electron transport chain defects in Alzheimer's disease brain. *Neurology* 44 (6):1090-1096
76. Mucke L, Masliah E, Yu GQ, Mallory M, Rockenstein EM, Tatsuno G, Hu K, Kholodenko D, Johnson-Wood K, McConlogue L (2000) High-level neuronal expression of abeta 1-42 in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J Neurosci* 20 (11):4050-4058
77. Butterworth RF, Besnard AM (1990) Thiamine-dependent enzyme changes in temporal cortex of patients with Alzheimer's disease. *Metab Brain Dis* 5 (4):179-184
78. Gibson GE, Haroutunian V, Zhang H, Park LC, Shi Q, Lesser M, Mohs RC, Sheu RK, Blass JP (2000) Mitochondrial damage in Alzheimer's disease varies with apolipoprotein E genotype. *Ann Neurol* 48 (3):297-303
79. Ko LW, Sheu KF, Thaler HT, Markesbery WR, Blass JP (2001) Selective loss of KGDHC-enriched neurons in Alzheimer temporal cortex: does mitochondrial variation contribute to selective vulnerability? *J Mol Neurosci* 17 (3):361-369. doi:10.1385/JMN:17:3:361
80. Mastrogiacoma F, Lindsay JG, Bettendorff L, Rice J, Kish SJ (1996) Brain protein and alpha-ketoglutarate dehydrogenase complex activity in Alzheimer's disease. *Ann Neurol* 39 (5):592-598. doi:10.1002/ana.410390508
81. Sorbi S, Bird ED, Blass JP (1983) Decreased pyruvate dehydrogenase complex activity in Huntington and Alzheimer brain. *Ann Neurol* 13 (1):72-78. doi:10.1002/ana.410130116
82. Casley CS, Land JM, Sharpe MA, Clark JB, Duchen MR, Canevari L (2002) Beta-amyloid fragment 25-35 causes mitochondrial dysfunction in primary cortical neurons. *Neurobiol Dis* 10 (3):258-267
83. Nunomura A, Perry G, Aliev G, Hirai K, Takeda A, Balraj EK, Jones PK, Ghanbari H, Wataya T, Shimohama S, Chiba S, Atwood CS, Petersen RB, Smith MA (2001) Oxidative damage is the earliest event in Alzheimer disease. *J Neuropathol Exp Neurol* 60 (8):759-767
84. Bogaert YE, Rosenthal RE, Fiskum G (1994) Postischemic inhibition of cerebral cortex pyruvate dehydrogenase. *Free Radic Biol Med* 16 (6):811-820
85. Zaidan E, Sheu KF, Sims NR (1998) The pyruvate dehydrogenase complex is partially inactivated during early recirculation following short-term forebrain ischemia in rats. *J Neurochem* 70 (1):233-241
86. Arsenian M (1998) Potential cardiovascular applications of glutamate, aspartate, and other amino acids. *Clin Cardiol* 21 (9):620-624
87. Wischmeyer PE, Jayakar D, Williams U, Singleton KD, Riehm J, Bacha EA, Jeevanandam V, Christians U, Serkova N (2003) Single dose of glutamine enhances myocardial tissue metabolism, glutathione content, and improves myocardial function after ischemia-reperfusion injury. *JPEN J Parenter Enteral Nutr* 27 (6):396-403
88. Jha MK, Jeon S, Suk K (2012) Pyruvate Dehydrogenase Kinases in the Nervous System: Their Principal Functions in Neuronal-glial Metabolic Interaction and Neuro-metabolic Disorders. *Curr Neuroparmacol* 10 (4):393-403. doi:10.2174/157015912804143586
89. Bielarczyk H, Gul S, Ronowska A, Bizon-Zygmanska D, Pawelczyk T, Szutowicz A (2006) RS-alpha-lipoic acid protects cholinergic cells against sodium nitroprusside and amyloid-beta neurotoxicity through restoration of acetyl-CoA level. *J Neurochem* 98 (4):1242-1251. doi:10.1111/j.1471-4159.2006.03966.x
90. Zhang L, Xing GQ, Barker JL, Chang Y, Maric D, Ma W, Li BS, Rubinow DR (2001) Alpha-lipoic acid protects rat cortical neurons against cell death induced by amyloid and hydrogen peroxide through the Akt signalling pathway. *Neurosci Lett* 312 (3):125-128
91. Hager K, Marahrens A, Kenkies M, Riederer P, Munch G (2001) Alpha-lipoic acid as a new treatment option for Alzheimer [corrected] type dementia. *Arch Gerontol Geriatr* 32 (3):275-282
92. Henderson ST, Poirier J (2011) Pharmacogenetic analysis of the effects of polymorphisms in APOE, IDE and IL1B on a ketone body based therapeutic on cognition in mild to moderate Alzheimer's disease; a randomized, double-blind, placebo-controlled study. *BMC Med Genet* 12:137. doi:10.1186/1471-2350-12-137

93. Reger MA, Henderson ST, Hale C, Cholerton B, Baker LD, Watson GS, Hyde K, Chapman D, Craft S (2004) Effects of beta-hydroxybutyrate on cognition in memory-impaired adults. *Neurobiol Aging* 25 (3):311-314. doi:10.1016/S0197-4580(03)00087-3

94. Vassar R, Kuhn PH, Haass C, Kennedy ME, Rajendran L, Wong PC, Lichtenthaler SF (2014) Function, therapeutic potential and cell biology of BACE proteases: current status and future prospects. *J Neurochem* 130 (1):4-28. doi:10.1111/jnc.12715

Figure legends

Figure 1: BACE1 overexpression alters APP processing and glucose oxidation in SH-SY5Y cells. **A.** Representative immunoblots showing stable overexpression of BACE1 and resultant changes in cellular APP and sAPP α and sAPP β shed into the culture media, compared to empty vector (EV) treated controls. **B. – E.** SH-SY5Y BACE1 overexpression did not change APP protein levels (**B**; n=4), increased total BACE1 protein levels (**C**; n = 7) and induced a shift in APP processing, with a reduction in sAPP α (**D**; n = 4) and an increase in sAPP β (**E**; n = 4). **F. – H.** SH-SY5Y BACE1 overexpression reduced ^{14}C -glucose oxidation rate (**F**; n = 7), with no reduction in ^{14}C -glucose incorporation rate (**G**; n = 9), giving an overall reduction in the ratio of ^{14}C -glucose oxidation to incorporation (**H**; n = 7). Values are means \pm SEM. ** $p < 0.01$, *** $p < 0.001$

Figure 2: Glucose oxidation in SH-SY5Y cells is not impaired following overexpression of secretase-dead BACE1 protein. **A.** Representative immunoblots showing stable overexpression of wild type BACE1 and a secretase-dead BACE1 mutant (mBACE1) and resultant changes in cellular APP and sAPP β shed into the culture media. **B. – D.** SH-SY5Y mBACE1 overexpression increased total BACE1 to levels equivalent to BACE1 overexpressed cells and ~3-4X greater than empty vector (EV) controls (**B**; n = 5), with no effect on APP levels (**C**; n = 3), but in contrast to BACE1 overexpression, mBACE1 reduced sAPP β to levels below that of the EV controls (**D**; n = 6 – 10). **E. – G.** SH-SY5Y mBACE1 overexpression had no effect on ^{14}C -glucose oxidation compared to EV controls, in contrast to the reduction in ^{14}C -glucose oxidation observed in BACE1 overexpressing cells (**E**; n = 5), with mBACE1 having no effect on ^{14}C -glucose incorporation (**F**; n = 5), resulting in the ratio of ^{14}C -glucose oxidation to incorporation being unchanged in mBACE1 cells (**G**; n = 5). Values are means \pm SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 3: BACE1 overexpression alters glucose metabolism in SH-SY5Y cells.

A. – C. BACE1 overexpression reduces oxygen consumption rate (OCR) (**A**) and increases the extracellular acidification rate (ECAR) (**B**) resulting in a reduced OCR:ECAR ratio (**C**), compared to EV cells (n = 7 – 12). **D.** Glycolysis stress test temporal profile for EV and BACE1 SH-SY5Y cells showing the effects on ECAR of the sequential addition of 2.5 mM glucose, 1 μM oligomycin (Oligo) and 25 mM 2-deoxyglucose (2DG). **E. – G.** BACE1 overexpression increased basal glycolysis (**E**), maximal glycolysis (**F**) and glycolytic reserve (**G**) compared to EV-treated cells (n = 18 – 21). Values are means \pm SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 4: BACE1 overexpression has no effect on ATP-linked, maximal or leak mitochondrial oxygen consumption.

A. Modified mitochondrial stress temporal profile for EV and BACE1 SH-SY5Y cells showing the effects on the percentage change in normalised OCR of the sequential addition of 1 μ M oligomycin (oligo) and 2 μ M rotenone and antimycin (Rot + Ant). **B. – D.** BACE1 overexpression had no effect on ATP-linked OCR (**B**), on the OCR required to maintain the mitochondrial leak (**C**) or the non-mitochondrial OCR (**D**) compared to EV cells (n = 8). **E.** Temporal profile showing the normalised baseline OCR for EV and BACE1 cells and the maximal OCR (reserve capacity) attained following addition of 0.2 μ M carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP). **F.** BACE1 overexpression has no effect on the percentage OCR increase following FCCP addition compared to EV cells (n = 9-10). Values are means \pm SEM.


Figure 5: BACE1 overexpression induces specific mitochondrial TCA cycle enzyme lesions. **A, B.** SH-SY5Y cells overexpressing BACE1 exhibit reduced lactate usage at levels \leq 2mM compared to EV cells as demonstrated by (**A**) reduced repression of 14 C-glucose oxidation (n = 6) and (**B**) direct OCR when lactate provided as sole substrate (n = 6 – 11). **C.** Pyruvate dehydrogenase activity (PDH) is reduced in BACE1 overexpressing, compared to EV, cells (n = 5). **D.** OCR when EV and BACE1 cells provided 2.5 mM pyruvate as sole substrate (n = 5). **E.** OCR of EV and BACE1 SH-SY5Y cells when provided with 2.5 mM glucose alone or 2.5 mM glucose + 4 mM glutamine (n = 4 – 10). **F.** direct OCR when 4 mM glutamine provided as sole substrate (n = 8). **G.** BACE1, but not mBACE1, overexpression, reduced the enzyme activity of α -KGDH, compared to EV cells (n = 6). **H.** IDH activity (NAD⁺ and NADP⁺ isoforms) is reduced by BACE1 overexpression compared to EV cells (n = 4). **I.** BACE1 or mBACE1 overexpression has no effect on fumarase enzyme activity, protein levels or specific activity compared to EV cells (n = 7). Values are means \pm SEM. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 6: Reversal of BACE1 mediated impaired glucose oxidation in SH-SY5Y cells.


A,B. Representative immunoblots (**A**) of phosphorylated PDHe1 α (p-PDHe1 α) subunit, total PDHe1 α and actin loading control in control (EV) and BACE1 overexpressing cells in the absence and presence of 10 μ M or 100 μ M dichloroacetate (DCA), with quantification of the immunoblot data shown in **B** (n = 4). **C.** DCA partially reverses BACE1 mediated impairment of glucose oxidation, but also increases OCR of EV cells (n = 5). **D.** reduced OCR associated with BACE1 cells is recovered to EV control levels by addition of 1 μ M and 10 μ M beta hydroxybutyrate (n = 4 – 6). **E.** Representative immunoblots of BCL-2-associated agonist of cell death (BAD) and actin loading control with quantification of the immunoblot data (n = 4). **F.** BACE1-mediated reduction in OCR is recovered to EV control levels by addition of 50 μ M α -lipoic acid (n = 6). Values are means \pm SEM. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 1.TIF


A




B




C




D




E



F



G



H




Figure 2.TIF




Figure 3.TIF




Figure 4.TIF




Figure 5.TIF




Figure 6.TIF

